

CO₂ EVOLUTION IN CAVE ATMOSPHERES: NEW INSIGHTS, CONCERNS AND MITIGATION STRATEGIES.

F. Bourges₁, F. Perrier₂, B. Lartiges₃, D.Genty₄, F. Girault₂, R. Losno₂, A. François₅, J. Leplat₅, S. Touron₅, F. Boustas₅, S. Tocino₆.

1) Géologie Environnement Conseil, 30 rue de la République, F-09200 Saint-Girons, France

2) Institut de Physique du Globe de Paris, Université de Paris, 1 rue Jussieu, F-75005 Paris, France

3) Université de Toulouse III Paul Sabatier, Géosciences Environnement-Toulouse, 14 av. Edouard Belin, F-31400 Toulouse, France

4) Environnements et Paléoenvironnements Océaniques et Continentaux (EPOC), UMR 5805, Université de Bordeaux, Allée Geoffroy Saint-Hilaire, F-33615 Pessac, France

5) Laboratoire de Recherches des Monuments Historiques (CRC, USR3224), 29 rue de Paris, F-77420 Champs-sur-Marne

6) Site de l'Aven d'Ornac, F-07150 Vallon-Pont d'Arc, France

This work is carried out by a research group : GIS-GEMS "Groupe d'Etude du Milieu Souterrain"

Motivation

CO₂ concentration upward drifts in cave atmospheres induce

- worrying levels of CO₂ concentration for human safety
- underground patrimony preservation

Study Sites

The Chauvet Cave has preserved a 360 centuries old rock art karstic massif of the French Massif central. The cave atmosphere is divided in two compartment with contrasted compositions on average (2.2% % CO₂ Vol. in the main volume (C1) and 3.4% in the rear room) C2) all are subject to smooth, 1.6 to 2% in amplitude, annual sinusoidal variations.

The Aven d'Ornac located at 7 km is a tourist-cave CO₂% with outstanding underground landscape and speleothems. Atmosphere composition variations are controlled by ventilation regimes. During the hot season, air is drained from the karst porosity (2.5 to 3 %Vol. CO₂ in the visited rooms, and 3 to 5% in remote networks). During the cold season, ventilation switches to a density driven regime: mixing with outside air falling from the upper entrance and dilutes the CO₂ concentration (<0.5%).




Methods

Long term multiparameter monitoring (since 1997) with continuous measurements and synchronized acquisition (15 mn time lag) were installed in the two caves and outside.




Natural context and evolution

Caves cross sections and ventilation patterns.




CO₂ long term evolutions

Chauvet Cave : a drift of + 0,5 % Vol. CO₂ is recorded in the confined Chauvet cave. Atmosphere in the cave is increasingly incompatible with human presence. Diverging climate situation can put the paintings at risk



Aven d'Ornac : The safety limit of 3% Vol. for visitors has been reached in 2016. Then forced ventilation is now necessary each year.





- Long term drifts of CO₂ in cave atmospheres can be attributed to a source effect : increase of vegetation density identified in most french karsts since 50 years.
- Seasonnal and pluriannual variations are a transfert effect mostly influenced by rainfall and subsequent infiltration in confined caves (Chauvet) and aerodynamic effect in open caves due to thermal effects (Ornac).

Remediation in the Aven d'Ornac

Stimulated ventilation that mimics the natural winter regime is operated in the visited part of the cave since 2016 safety alert.

Air pumping by the existing lift shaft is optimized to dilute CO₂ and radon and to keep characteristics of a subterranean atmosphere due to natural air input flux from the porosity of the karst volume.



Lessons

- The atmosphere of many caves evolves significantly (+ 5000 ppm in 23 years in Chauvet) due to constraints of the local environment and climate change consequences.
- In touristic caves as in l'Aven d'Ornac, stimulated ventilation is used to dilute CO₂. It mimics a natural regime and keeps the characteristic features of the cave atmosphere.
- In fragile painted caves, direct action on cave aerology is not recommended and only actions on the CO₂ source driver (vegetation cover) appears to be possible.

References

Hydrogeological control on carbon dioxide input into the atmosphere of the Chauvet-Pont d'Arc cave. F. Bourges, D. Genty F. Perrier, B. Lartiges, É. Régner, A. François, J. Leplat, S. Touron, F. Boustas, M. Massault, M. Delmotte, J.-P. Dumoulin, F. Girault. M. Ramonet, Ch. Chauveau, P. Rodrigues, Science of The Total Environment, Vol. 716, 10 May 2020.

Conservation of prehistoric caves and stability of their inner climate: lessons from Chauvet and other French caves. Bourges F. , Genthon P., Genty D. , Lorblanchet M. , Mauduit E. , D'Hulst D. Science of the Total Environment. Vol. 493, 15 Sept. 2014, p. 79-91DOI:10.1016/j.scitotenv.2014.05.137

Microclimates of l'Aven d'Ornac and other French limestone caves (Chauvet, Esparros, Marsoulas). BOURGES François, GENTHON Pierre, MANGIN Alain and D'HULST Dominique. International Journal of Climatology, 2006, 26 (12), p. 1651-1670. Wiley InterScience.